

The Ant Fauna of the Oriental Region: An Overview (Hymenoptera, Formicidae)¹⁾

Kazuo Ogata²⁾

Abstract

The number of species of ants in the Oriental region are estimated more than 3,000, belonging to 135 genera of 9 subfamilies. A list of genera in the Oriental region is presented. At genus level, 34 of them are endemic to this region. The Oriental and the Australian regions share 82 genera in common, which represents about 49 % of total genera in both regions. This is the highest ratio of similarity comparing with other regions.

Current taxonomic works try to revise higher taxa based on extensive material. As a result, the number of genera has been reducing due to synonyms, but that of species has been increasing. In spite of the effort, there are still many genera left unstudied, so that 30 to 80 % of species collected from certain area cannot be identified. The present confusion comes from (1) isolate descriptions without appropriate comparison; (2) obscure definition of higher taxa; (3) unreliable "intraspecific variations". A good network of researchers will help to solve the present confusion.

Introduction

The present knowledge on the ant fauna in the Oriental region is quite obscure. The information on distribution of each ant in the region is incomplete and fragmental. According to Wilson (1987), approximately 8,800 species belonging to 303 genera of 11 subfamilies are described from the world to that time. The pie graph of Fig. 1, drawn from his data, shows one-fourth of the total number of species occurs in Asia. His estimation, however, leaves two points untouched. Firstly, the cited data in his estimation were not the same level of accuracy. As for the Asian ants, the figure was cited from the check list by Chapman & Capco (1951). This literature has been the only one covering this region but outdated. Indeed, many taxonomic changes and additions of new taxa have been made since that time. Thus the figure does not reflect the actual number. Secondly, this is not the estimation of the Oriental region.

1) This paper is a modification of the poster presented at the XIXth International Congress of Entomology, Beijing, China, 1992.

2) Institute of Tropical Agriculture, Kyushu University, Fukuoka 812, Japan.




Fig. 1. Relative distribution of estimated numbers of species in the world (after Wilson, 1987).

In this paper, I would like to show the present situation of the ant taxonomy in the Oriental region. To begin with, the statistics of genera in the Oriental region is given with a discussion of biogeographical characteristics of the region. Then, a brief history of taxonomic activities and current state of the classification are overviewed. Finally, a prospect of ant taxonomy in this region is discussed.

The boundary of the Oriental region here follows tentatively that of Brown (1973). That is, the region consists of Indian subcontinent including Pakistan, south of the Pamir and the Himalayas, Myanmar (Burma), Indochinese and Malay Peninsulas, Indonesia including Malay Archipelago (the East Indies) to Timor and Sulawesi (Celebes), the Philippines, Taiwan, southern China from the Tsinling Mountains and the Tibetan Scarp, and southern Japan.

The subdivisions of the Oriental region are provisionally adopted from Lincoln & Boxshall (1987). The following four subregions are distinguished: the Indian subregion covering base of the Indian Peninsula; the Ceylonese subregion covering southern India and Sri Lanka; the Indo-Chinese subregion covering south of the Himalayas, Myanmar, Indochina, southern China, Taiwan and south of Japan; the Indo-Malayan subregion covering the Malay Peninsula, and SE Asian islands west of Weber's line. I admit that a detailed analysis for the exact limitation of the region and its subdivisions are necessary. This must be one of the future subject.

Genera of the Oriental region

At present, 135 genera of 9 subfamilies are recognized in the Oriental region. These are listed in the appendix. Since there have been no comprehensive check lists of this region yet, the exact number of species is still obscure, but would be estimated over 3,000.

A comparison of the number of genera among 7 zoogeographical regions is shown in Table 1. At genus level, the Oriental region has the highest number. Although several genera are questionable for their taxonomic status and they might be submerged as synonyms, it is supposed that there are still undescribed genera. This is supported by the fact that even within recent 5 years 12 genera have been described as new.

In terms of the similarity of generic composition, the Australian regions is the closest. The number of common genera in the Oriental and Australian regions is 82, which corresponds to 48.5 % of total number in both regions. On the other hand, the Palearctic region has a lower ratio of similarity (28.9 %), in spite of its close geographical position. Brown (1973) stated that the Oriental and Australian regions are not separated by a cold-temperature barrier and thus "the Oriental merges gradually into the Australian region within the tropics". Temperature factor would be applicable for a lower similarity between the Oriental and Palearctic regions, though the transition zone is rather broad in the boundary between these regions in their eastern part. An analysis of historical factor is incomplete because phylogenetic information is quite inadequate.

Table 1. Number of endemic genera in each zoogeographical region and that of common genera with the Oriental region. AU, Australian; ET, Ethiopian; MA, Malagasy; NA, Nearctic; NT, Neotropical; OR, Oriental; PA, Palearctic. Dash indicates absence of the subfamily. 0 means no endemic genera in the region.

	OR	AU	PA	ET	MA	NA	NT
Nothomyrmecinae	—	1	—	—	—	—	—
Myrmecinae	—	1	—	—	—	—	—
Ponerinae	1	1	1	12	1	0	7
Cerapachyinae	0	0	0	0	0	1	3
Dorylinae	0	0	0	0	—	—	—
Ecitoninae	—	—	—	—	—	0	2
Leptanillinae	5	0	0	1	—	—	0
Myrmicinae	19	10	8	11	3	1	35
Pseudomyrmecinae	0	0	—	0	0	0	1
Aneuretinae	1	—	—	—	—	—	—
Dolichoderinae	1	4	0	2	0	0	2
Formicinae	7	9	2	4	0	2	4
No. of endemic genu.	34	26	11	30	4	4	54
No. of genu.	135	116	61	96	37	69	123
% of endemic genu.	25.2	22.4	18.0	31.3	10.8	5.8	43.9
Common genu. with OR	—	82	44	61	30	38	44
Common Index*	—	48.5	28.9	35.9	21.1	22.8	20.7

* Common Index = $\frac{\text{no. of genu. in common}}{\text{total no. of genu. in both areas}} \times 100$

Table 2. List of endemic genera to the Oriental region. O1, Indian subregion; O2, Ceylonese subregion; O3, Indo-Chinese subregion; O4, Indo-Malayan subregion. Asterisks show informal estimation based on personal communications with R.W. Taylor (on *Anomalomyrma* and *Protanilla*) and with M. Terayama (on *Trigonogaster*).

	spp.	Distribution					spp.	Distribution			
		O1	O2	O3	O4			O1	O2	O3	O4
PONERINAE						<i>Pentastruma</i>	2	-	-	+	-
<i>Harpegnathos</i>	5	-	+	+	+	<i>Proatta</i>	1	-	-	-	+
LEPTANILLINAE						<i>Rotastruma</i>	2	-	-	+	+
<i>Anomalomyrma</i>	*>2	-	-	+	+	<i>Secostruma</i>	1	-	-	-	+
<i>Noonilla</i>	1	-	-	-	+	<i>Stereomyrmex</i>	1	-	+	-	+
<i>Phaulomyrma</i>	1	-	-	-	+	<i>Tetheamyrmia</i>	1	-	-	-	+
<i>Protanilla</i>	*>2	+	+	+	+	<i>Tricyclurus</i>	1	-	-	-	+
<i>Scyphodon</i>	1	-	-	-	+	<i>Trigonogaster</i>	*>2	+	+	+	+
MYRMICINAE						ANEURETINAE					
<i>Acanthomyrmex</i>	11	-	+	+	+	<i>Aneuretus</i>	1	-	+	-	-
<i>Anillomyrma</i>	2	-	+	-	+	DOLICHODERINAE					
<i>Asketogenys</i>	1	-	-	-	+	<i>Loweriella</i>	1	-	-	-	+
<i>Chimaeridris</i>	2	-	-	-	+	FORMICINAE					
<i>Dysedrognathus</i>	1	-	-	-	+	<i>Bregmatomyrma</i>	1	-	-	-	+
<i>Epelysidris</i>	1	-	-	-	+	<i>Cladomyrma</i>	5	-	-	-	+
<i>Indomyrma</i>	1	-	+	-	-	<i>Forelophilus</i>	1	-	-	-	+
<i>Ishakidris</i>	1	-	-	-	+	<i>Gesomyrmex</i>	6	-	-	+	+
<i>Kartidris</i>	2	+	-	+	-	<i>Myrmoteras</i>	18	-	+	-	+
<i>Lophomyrmex</i>	?4	+	-	+	+	<i>Overbeckia</i>	1	-	-	-	+
<i>Paratpula</i>	9	-	+	-	+	<i>Pseudophomomyrmex</i>	1	-	-	-	+

Endemic genera

Brown (1973) estimated 22 endemic genera in the Oriental region and referred to as the lowest endemism among the tropical regions. Now the genera are counted 34 (Table 2). This increase is mainly because of newly established genera, in addition to taxonomic changes, re-examinations of distributional data. Indeed 11 genera have been described since 1973. As a result, the number is comparable to that in the Ethiopian region. But the percentage of endemic genera is not so higher as in two other tropical regions, the Ethiopian and Neotropical regions. The ratio is almost same as in the Australian region.

The lower ratio of endemism might concern with the boundary between Sulawesi and New Guinea. If, ignoring the boundary, we consider the tropical and subtropical areas of the Oriental-Australian region, the ratio would become comparable with other two tropical regions.

Distribution patterns of endemic genera are shown in Table 2. Among 34 endemic genera, 2 genera, *Protanilla* and *Trigonogaster*, are distributed through the whole region. The Indo-Malayan subregion is most abundant in endemic genera. As many as 18 genera occur in this subregion exclusively, though several of them are seemed to be question in their taxonomic position. Other genera having more or less restricted distribution are *Indomyrma*,




Fig. 2. Distribution patterns of some endemic genera to the Oriental region. a, *Harpegnathos*; b, *Myrmoterus*; c, *Aneuretus* and *Pentastruma*.

Pentastruma (Fig. 2c), *Stereomyrmex*, and *Aneuretus* (Fig. 2c). They are found in the Ceylonese subregion or Indo-Chinese subregion. No genera occur in the Indian subregion exclusively.

The distributions of other endemic genera are divided into three patterns. The first one includes the genera found mainly in the Indo-Chinese subregion with adjacent areas. Examples of this pattern are *Harpegnathos* (Fig. 2a), *Kartidris* and *Lophomyrmex*.

The second pattern is shown by the genera found mainly in southern India and SE Asian islands (O2+O4). This pattern is represented by *Anillomyrma*, *Acanthomyrmex*, *Paratpula* and *Myrmoterus* (Fig. 2b).

The third pattern includes the genera *Anomalomyrma*, *Rotastruma* and *Gesomyrmex* which are found in eastern areas (O3+O4).

Genera occurred in the Oriental and other tropical regions

The genera only found both in the Oriental and Australian regions are many (21 of 82 shared genera occur in the both regions exclusively). As represented by the pattern of *Myopias* (Fig. 3b), many of them are found in a belt from southwestern India to northeastern




Fig. 3. Distribution patterns of some genera occurring in the Oriental and other tropical regions.
a, *Cataulacus*; b, *Myopias*; c, *Eurhopalothrix*.

Australia. In particular, New Guinean ant fauna is important in considering the biogeography of both regions.

The genera found both in the Ethiopian (including the Malagasy) and the Oriental regions exclusively are counted 3: *Cataulacus* (Fig. 3a), *Myrmicaria*, *Paedalgus*, all of which belong to subfamily Myrmicinae but have phylogenetically no close relationships. The distributions of these genera show following two common patterns: 1) at least they occur in the Ceylonese subregion; 2) the species number is fewer to the eastward of the Oriental region. In terms of species number for these 3 genera, the Ethiopian region has more species than the Oriental region (46 : 17 in *Cataulacus*; 3 : 1 in *Paedalgus*) or at most the both have almost same number (9 : 8 in *Myrmicaria*), though the actual number of species might be increased because of the lack of recent revisionary works except for *Cataulacus* (Bolton 1974, 1982).

There are only two genera distributed both in the Oriental and Neotropical regions exclusively: *Belonopelta* of Ponerinae and *Perissomyrmex* of Myrmicinae. The species of the *Belonopelta* in the Oriental region were formerly treated under the genus *Emeryopone* which was synonymized by Baroni Urbani (1975). *Perissomyrmex* is said to be of the Oriental origin (Hölldobler & Wilson, 1990). Taking into the Australian region, a distribution pattern like *Eurhopalothrix* (Fig. 3c) would suggest a Gondwanic origin.

Detailed analysis of phylogenys among species is awaited, which will lead to hypotheses of dispersal/vicariance history on faunal relationships.

Current State of Classification

Taxonomic changes

Over the past few decades a considerable number of studies have been made on the taxonomy of the Oriental ants. To see the tendency of such taxonomic activities, we will make several historical comparisons.

Table 3 indicates the number of genera in Chapman & Capco (1951) and that in 1991. Though the statistics is for Asian ants, it is almost same tendency for the Oriental ants. Apparently the total number of genera in 1991 is reduced from that in 1951. During 40 years, 34 genera have been added as new, status raised, or new records. On the other hand, reduction of number mainly due to synonyms is counted 51, exceeding the number of additions. In particular the decrease is distinct in the subfamily Ponerinae. This is because the taxon has been intensively revised. The reclassification of this subfamily has been greatly contributed by Brown (1958, 1960, 1975, 1978, etc.). This tendency is also true for the subfamily Dolichoderinae (see the appendix where the number of genera decreased to 9), though the Table does not show the latest figure since the revisions were made quite recently (Shattuck, 1992a, b, c). In any case, once a taxonomic review is made, the number is tend to reduced. Is it true for species-level taxonomy?

A historical review of classification of the tribe Tetramoriini reveals the tendency of taxonomy more clearly. Fig. 4 shows changes in the numbers of taxa of the tribe in the Oriental region.

In 1922 when the framework of higher classification was established by Emery in *Genera*

Table 3. Comparison of number of Asian genera between 1951 and 1991. A1, newly described; A2, status raised; A3, new record; R1, synonyms; R2, removals due to incorrect locality; S, subfamily change.

	1951	addition			reduction		S	1991
		A1	A2	A3	R1	R2		
Ponerinae	37	0	5	0	16	1	-1	24
Cerapachyinae	4	0	0	1	3	0	+1	3
Leptanillinae	2	5	0	0	0	0	0	7
Dorylinae	2	0	0	0	0	0	0	2
Pseudomyrmecinae	1	0	0	0	0	0	0	1
Myrmicinae	71	18	0	2	19	2	0	70
Aneuretinae	—	0	0	0	0	0	+1	1
Dolichoderinae	14	0	2	1	3	0	-1	13
Formicinae	29	0	0	0	7	1	0	21
Total	160	23	7	4	48	4	—	142




Fig. 4. Historical changes in the number of taxa of the tribe Tetramoriini in the Oriental region. Aci, *Acidomyrmex*; Dyo, *Dyomorium*; Ire, *Ireneela*; Iip, *Ireneopone*; Rho, *Rhoptromyrmex*; Sec, *Secostruma*; Tet, *Tetramorium*; Tri, *Triglyphothrix*; Xip, *Xiphomyrmex*. Unshaded areas show the number of infraspecific taxa.

Insectorum, the tribe includes 4 genera in the Oriental region. At that time 79 species and infraspecific forms had been described.

Almost thirty years after, by the publication of the check list of Chapman & Capco (1951), 4 genera had been added: *Acidomyrmex* raised from a subgenus of *Rhoptromyrmex*, *Dyomorium* Donisthorpe, *Ireneela* Donisthorpe, and *Ireneopone* Donisthorpe. To that time, the tribe includes 88 species and infraspecific forms. It should be noted that the increase of taxa mainly attribute to infraspecific ones. In this duration, many taxonomic papers were presented but most of them were isolated descriptions.

Since the 1950's, revisionary works based on extensive material and detailed type comparisons have been increased. As for the classification of Tetramoriini, Brown (1964) revised *Rhoptromyrmex*, where *Acidomyrmex* was synonymized with it. In his revision of Solenopsidini and Pheidologetini, Ettershank (1966) synonymized *Dyomorium* with *Vollenhovia* (outside of the tribe). Most of the taxonomic changes in this duration were made by Bolton (1976, 1977). As a result, by 1981, the number of species-group taxa in the tribe was increased to 102, but that of genera became only 3.

Bolton further added *Triglyphothrix* into the synonymy of *Tetramorium* in 1985 and described a monotypic genus *Secostruma* as new in 1986. Presently we have 3 genera comprising over 100 species of the tribe Tetramoriini.

From these examples, we can see the tendency of taxonomic changes as follows:

- (1) The number of genera has been decreasing due to synonyms.




Fig. 5. Relative numbers of unidentified materials in various areas.

- (2) The number of species has been increasing by finding new species.
- (3) Intraspecific forms has been disappearing.

In addition to these taxonomic activities, the current studies of taxonomy also tend to focus on the phylogeny of higher taxa (e.g. Bolton, 1990b, 1990c; Ogata, 1990; Shattuck, 1992b; Ward, 1990).

Limit of identification

That is the case where the taxonomic revisions have been made recently. But such genera are limited (recent revisionary works are shown in the appendix). Most genera in the Oriental region have been left unstudied. Take several faunal reports for examples (Fig. 5).

In a series of karyological survey, Imai and his colleagues tried to identify their specimens exactly as possible. Despite the collaboration with American or European authorities of taxonomy, their material has high ratio of unidentified species (about 30 to 80 %), even it might include undescribed ones.

The situation is also true for the Nansei Islands of Japan. The islands are situated between Taiwan and Kyushu of Japan and are transition zone from the Oriental region to the Palearctic region in species composition. We can distinguished 170 species, but about 40 % of them are taxonomically unsolved (Ogata, unpublished).

The present situation of ant taxonomy in the Oriental region is simply expressed by the word "Oriental Chaos" (Brown, pers. com). Taxonomic studies before 1950's concentrate on

descriptions of local faunas. Have such efforts contributed the development of ant taxonomy? Brown (1955) stated the history as follows:

"The years following 1910 saw many specialists joined the rush to describe ants... Taken generally, their work is very disappointing...."

Indeed it is often difficult to identify the species without reference to the types because most of the descriptions and keys are poor and useless. Furthermore, finding the type materials itself is sometimes difficult because they are scattered in various institutes of American or European countries. As seen in the previous section, the confusion comes from: (1) isolate descriptions without appropriate comparison; (2) obscure definition of higher taxa; (3) unreliable "infra-specific variations".

This state is not restricted to the ants of the Oriental region (see Taylor, 1983 for Australia), but unlike the Australian National Insect Collection there are few institutes having comprehensive collections in Asian countries.

Prospect

Wilson (1987) supposed that 20,000 or more species of ants exist in the world, and that the number of genera will reach about 350. The survey and synthesis of the Oriental region are on going by many taxonomists. Year by year, new knowledge is adding. But since the family Formicidae is such a huge group, it would need time to construct a systematized information. On the other hand, in a recent crisis of biodiversity, describing a local fauna, in particular that in the tropics, becomes a urgent and important subject (Wilson, 1988). But considering the previous history which caused the present confusion, we should not only concentrate on a local fauna but have comprehensive view. Then, is there any effective way to solve the present situation?

As illustrated by Taylor (1983), there are several levels between species in nature and species taxonomically understood. To increase species taxonomically understood, a rate of taxonomic activity should be advanced. Information networks will play an important role in this rate. In case of Japan, the Myrmecological Society of Japan have published a series of identification manuals (1988-1992). The books cover all the distinguishable species, including ones not formally named or "problematical species". This is tentative treatment, but we can distinguish the species taxonomically understood from ones in question which will be treated in future revisionary works with extensive material and accumulated information. In carrying the project, the networks of myrmecologists has been quite useful and now the knowledge on the taxonomy of Japanese ants is held in common by more researchers at least in Japan.

The Oriental region, like other tropical regions, consists of many developing countries. This means that there are still many areas where scientific survey is difficult due to political

reasons or ill transportation system. Using a good networks of world researchers can avoid further confusion and accelerate the rate of taxonomic activity.

Acknowledgements

I wish to thank the following persons for their constant guidance and encouragement: W. L. Brown, Jr., R. W. Taylor, M. Terayama, M. Kubota, H. T. Imai, K. Masuko. This study was supported in part by a grant from the Nippon Life Insurance Foundation.

Appendix: List of genera in the Oriental region

This list is based on a table by Hölldobler & Wilson (1990) with a modification of later taxonomic changes. Genera are listed alphabetically within each subfamily. New additions since Chapman & Capco's check list (1951) are shown by asterisks. Numbers are put in presently recognizable names in the Oriental regions. Bold face genera show the endemic to the Oriental region. Italics show junior synonyms or removals from the region. Arrows indicate correct regions where the excluded genus is occurred. Abbreviations of zoogeographical regions are as in Table 1 (small letters of those indicate the region introduced).

PONERINAE

- | | | |
|--------------------------------------|---------------------------|----------------------|
| 1. Amblyopone Erichson 1842 | AU, ET, NA, NT, OR,
PA | Brown, 1960 |
| 2. Anochetus Mayr 1861 | AU, ET, NT, OR | Brown, 1978 |
| *3. Belonopelta Mayr 1870 | NT, OR | Baroni Urbani, 1975a |
| <i>Bothroponera</i> Mayr 1862 | =Pachycondyla | |
| *4. Brachyponera Emery 1901 | AU, ET, OR | |
| 5. Centromyrmex Mayr 186 | ET, NT, OR | |
| <i>Chalcoponera</i> Emery 1897 | =Rhytidoponera | |
| 6. Cryptopone Emery 1892 | AU, ET, NA, NT, OR,
PA | |
| 7. Diacamma Mayr 1862 | AU, OR | |
| 8. Discothyrea Roger 1863 | AU, ET, NA, NT, OR | Brown, 1958 |
| 9. Ectomomyrmex Mayr 1867 | AU, OR | |
| <i>Emeryopone</i> Forel 1912 | =Belonopelta | |
| <i>Euponera</i> Forel 1891 | →[MA] | |
| <i>Examblyopone</i> Donisthopre 1949 | =Prionopelta | |
| *10. Gnampotogenys Roger 1863 | AU, NA, NT, OR | Brown, 1958 |

- | | | | |
|------|--------------------------------------------|-------------------------------|----------------------------|
| 11. | Harpegnathos Jerdon 1851 | OR | |
| *12. | Hypoponera Santschi 1938 | AU, ET, MA, NA, NT,
OR, PA | Taylor, 1967 |
| 13. | Leptogenys Roger 1861 | AU, ET, MA, NA, NT,
OR, PA | |
| | <i>Microbolbos</i> Donisthorpe 1948 | =Leptogenys | |
| 14. | Myopias Roger 1861 | AU, OR | |
| 15. | Myopopone Roger 1861 | AU, OR | Brown, 1960 |
| 16. | Mystrium Roger 1862 | AU, ET, MA, OR | Brown, 1960 |
| 17. | Odontomachus Latreille 1804 | AU, ET, MA, NA, NT,
OR, PA | Brown, 1976 |
| 18. | Odontoponera Mayr 1862 | AU, OR | |
| 19. | Pachycondyla F.Smith 1858 | AU, ET, MA, NA, NT,
OR, PA | [incl. Bothroponera] |
| 20. | Platythyrea Roger 1863 | AU, ET, NA, NT, OR | Brown, 1975 |
| 21. | Ponera Latreille 1804 | AU, ET, NA, NT, OR,
PA | Taylor, 1967 |
| 22. | Prionopelta Mayr 1866 | AU, ET, MA, NA, NT,
OR | Brown, 1960 |
| 23. | Probolomyrmex Mayr 1901 | AU, ET, NT, OR | Brown, 1975 |
| 24. | Proceratium Roger 1863 | AU, ET, MA, NA, NT,
OR, PA | Brown, 1958 |
| | <i>Prodiscothyrea</i> Wheeler 1916 | =Discothyrea | |
| | <i>Pseudocryptopone</i> Wheeler 1933 | =Ponera | |
| | <i>Pseudoneoponera</i> Donisthorpe
1943 | =Pachycondyla | |
| | <i>Pseudoponera</i> Emery 1901 | =Pachycondyla | |
| | <i>Renea</i> Donisthorpe 1947 | =Prionopelta | |
| | <i>Rhopalopone</i> Emery 1897 | =Gnamptogenys | |
| 25. | Rhytidoponera Mayr 1862 | AU, OR | Brown, 1958; Ward,
1980 |
| | <i>Selenopone</i> Wheeler 1933 | =Ponera | |
| | <i>Stictoponera</i> Mayr 1887 | =Gnamptogenys | |
| | <i>Stigmatomma</i> Roger 1859 | =Amblyopone | |
| | <i>Sysphincta</i> Roger 1863 | =Proceratium | |
| 26. | Trachymesopus Emery 1911 | AU, ET, OR | |
| | <i>Trapeziopelta</i> Mayr 1862 | =Myopias | |

CERAPACHYINAE [For the status of the subfamily and phylogenetic relationship, see Bolton (1990a, c)]

- | | | | |
|------|-------------------------------|-------------------------------|-------------|
| 27. | Cerapachys F.Smith 1857 | AU, ET, MA, NA, NT,
OR, PA | Brown, 1975 |
| | <i>Eusphinctus</i> Emery 1893 | =Sphinctomyrmex | |
| | <i>Lioponera</i> Mayr 1878 | =Cerapachys | |
| | <i>Phyracaces</i> Emery 1902 | =Cerapachys | |
| 28. | Simopone Forel 1891 | AU, ET, MA, OR | Brown, 1975 |
| *29. | Sphinctomyrmex Mayr 1866 | AU, ET, NT, OR | Brown, 1975 |

LEPTANILLINAE [For the higher classification of the subfamily, see Bolton (1990b)]

- | | | | |
|------|----------------------------------------------|----------------|----------------------|
| *30. | Anomalomyrma Taylor in Bolton | OR | |
| | 1990 | | |
| 31. | Leptanilla Emery 1870 | AU, ET, OR, PA | Baroni Urbani, 1977c |
| *32. | Noonilla Petersen 1963 | OR | Baroni Urbani, 1977c |
| 33. | Phaulomyrma Wheeler &
Wheeler 1930 | OR | Baroni Urbani, 1977c |
| *34. | Protanilla Taylor in Bolton 1990 | OR | |
| *35. | Scyphodon Brues 1925 | OR | Baroni Urbani, 1977c |
| *36. | Yavnella Kugler 1986 | OR, PA | |

DORYLINAE [Bolton (1990c) established the subfamily Aenictinae comprising sole genus *Aenictus*]

- | | | | |
|-----|------------------------|------------|--------------|
| 37. | Aenictus Shuckard 1840 | AU, ET, OR | Wilson, 1964 |
| 38. | Dorylus Fabricius 1793 | ET, OR, PA | Wilson, 1964 |

PSEUDOMYRMECINAE

- | | | | |
|-----|-------------------------|----------------|------------|
| 39. | Tetraopone F.Smith 1852 | AU, MA, OR, PA | Ward, 1990 |
|-----|-------------------------|----------------|------------|

MYRMICINAE

- | | | | |
|-----|----------------------------------|---------------------------|---------------|
| | <i>Acalama</i> M. Smith 1948 | =Vollenhovia | |
| 40. | Acanthomyrmex Emery 1892 | OR | Moffett, 1986 |
| | <i>Acidomyrmex</i> Emery 1915 | =Rhoptromyrmex | |
| 41. | Adelomyrmex Emery 1897 | AU, NT, OR | |
| | <i>Aeromyrma</i> Forel 1891 | =Oligomyrmex | |
| | <i>Amauromyrmex</i> Wheeler 1929 | =Pheidologeton | |
| | <i>Ancyridis</i> Wheeler 1935 | → [AU: PNG] | |
| | <i>Aneleus</i> Emery 1900 | =Oligomyrmex | |
| 42. | Anillomyrma Emery 1913 | OR | Bolton, 1987 |
| 43. | Aphaenogaster Mayr 1853 | AU, MA, NA, NT, OR,
PA | |

- Aratromyrmex* Stitz 1938 =Liomyrmex
- *44. **Asketogenys** Brown, 1972 OR
- Atopula* Emery 1912 =Tetramorium
- Atta* Fabricius 1804 → [NA, NT]
45. *Calyptomyrmex* Emery 1887 AU, ET, OR Baroni Urbani, 1975b;
Bolton, 1981
46. *Cardiocondyla* Emery 1869 AU, ET, MA, na, nt,
OR, PA Bolton, 1982
47. *Carebara* Westwood 1894 ET, NT, OR
48. *Cataulacus* F.Smith 1853 ET, MA, OR Bolton, 1974, 1982
- Ceratopheidole* Pergande 1859 =Pheidole
- *49. **Chimaeridris** Wilson 1989 OR
- Conothorax* Karawajew 1935 =Pheidole
50. *Crematogaster* Lund 1831 AU, ET, MA, NA, NT,
OR, PA
- *51. *Dacatinops* Brown & Wilson 1957 AU, OR Taylor, 1985
52. *Dilobocondyla* Santschi 1910 AU, OR
- Dodous* Donisthorpe 1946 =Pristomyrmex
- Dyomorium* Donisthorpe 1947 =Vollenhovia
- *53. **Dysedrognathus** Taylor 1968 OR
- Eneria* Donisthorpe 1948 =Strumigenys
- *54. **Epelysidris** Bolton 1987 OR
55. *Epitritus* Emery 1869 ET, OR, PA
- Epopostruma* Forel 1865 → [AU]
- *56. *Eurhopalothrix* Brown & Kempf 1961 AU, NA, NT, OR Taylor, 1990a
- Gauromyrmex* Menozzi 1933 =Vollenhovia
- *57. *Glomyrmex* Wheeler 1915 AU, ET, NT, OR
- *58. **Indomyrma** Brown 1985 OR
- Ireneela* Donisthorpe 1941 =Rhoptromyrmex
- Ireneidris* Donisthorpe 1943 =Monomorium
- Ireneopone* Donisthorpe 1946 → [MA]
- Ischnomyrmex* Mayr 1862 =Pheidole
- *59. **Ishakidris** Bolton 1984 OR
- *60. **Kartidris** Bolton 1991 OR
- *61. *Kyidris* Brown 1949 AU, OR Wilson & Brown, 1956
62. *Leptothorax* Mayr 1855 AU, ET, MA, NA, NT,
OR, PA Bolton, 1982
63. *Liomyrmex* Mayr 1865 AU, OR

- | | | | |
|------|------------------------------------------------------------------|-------------------------------------------|-----------------------------------|
| 64. | Lophomyrmex Emery 1892 | OR | |
| 65. | Lordomyrma Emery 1897 | AU, OR | |
| *66. | Mayriella Forel 1902 | AU, OR | Baroni Urbani, 1977a |
| 67. | Meranoplus F.Smith 1853 | AU, ET, MA, OR | Bolton, 1981 |
| 68. | Messor Forel 1890 | ET, NA, OR, PA | Bolton, 1982 |
| 69. | Metapone Forel 1900 | AU, MA, OR | |
| 70. | Monomorium Mayr 1855 | AU, ET, MA, NA, NT,
OR | Ettershank, 1966;
Bolton, 1987 |
| 71. | Myrmecina Curtis 1829 | AU, NA, NT, OR, PA | |
| 72. | Myrmica Latreille 1804 | NA, OR, PA | |
| 73. | Myrmecaria Saunders 1841 | ET, OR | |
| 74. | Oligomyrmex Mayr 1867
<i>Orectognathus</i> F. Smith 1853 | AU, ET, MA, OR, PA
→ [AU] | Ettershank, 1966 |
| 75. | Paedalgus Forel 1911 | ET, OR | |
| 76. | Paratopula Wheeler 1919 | OR | Bolton, 1988a |
| 77. | Pentastruma Forel 1912 | OR | Brown & Boisvert,
1978 |
| *78. | Perissomyrmex M.Smith 1947 | NT, OR | |
| 79. | Pheidole Westwood 1840 | AU, ET, MA, NA, NT,
OR, PA | |
| 80. | Pheidologeton Mayr 1862 | AU, ET, OR | Ettershank, 1966 |
| 81. | Podomyrma F.Smith 1859 | AU, OR | |
| 82. | Pristomyrmex Mayr 1866 | AU, ET, MA, OR | Taylor, 1965; Bolton,
1981 |
| 83. | Proatta Forel 1912 | OR | |
| *84. | Quadristruma Brown 1949 | AU, ET, or | Bolton, 1983 |
| 85. | Rhopalomastix Forel 1900
<i>Rhopalothrix</i> Mayr 1870 | AU, OR
→ [AU, NT] | Taylor, 1991 |
| 86. | Rhopotromyrmex Mayr 1901
<i>Rogeria</i> Emery 1894 | AU, ET, OR
→ [AU, NA, NT] | Bolton, 1986 |
| 87. | Romblonella Wheeler 1935 | AU, OR | Taylor, 1990b |
| *88. | Rotastruma Bolton 1991 | OR | |
| *89. | Secostruma Bolton 1988 | OR | |
| 90. | Smithistruma Brown 1948

<i>Solenomyrma</i> Karawajew 1935 | ET, MA, NA, NT, OR,
PA
=Vollenhovia | |
| 91. | Solenopsis Westwood 1840 | AU, ET, NA, NT OR,
PA | Ettershank, 1966;
Bolton, 1987 |
| 92. | Stenamma Westwood 1840 | NA, NT, OR, PA | Baroni Urbani, 1977b |

93. **Stereomyrmex** Emery 1901 OR
94. *Strumigenys* F.Smith 1860 AU, ET, MA, NA, NT,
OR, PA
- *95. **Tetheamyrmex** Bolton, 1991 OR
96. *Tetramorium* Mayr 1855 AU, ET, MA, NA, NT, Bolton, 1977
OR, PA
Trichomyrmex Mayr 1865 =Monomorium
- *97. *Trichoscapa* Emery 1869 ET, na, nt, OR, PA Bolton, 1983
98. **Tricytarus** Donisthorpe 1947 OR
Triglyphothrix Forel 1890 =Tetramorium
99. **Trigonogaster** Forel 1890 OR
100. *Vollenhovia* Mayr 1865 AU, ET, na, OR, PA
- *101. *Vombisidris* Bolton, 1991 AU, OR
Wheeleriella Forel 1905 =Monomorium
Xiphomyrmex Forel 1887 =Tetramorium

ANEURETINAE

102. **Aneuretus** Emery 1892 OR Shattuck, 1992a

DOLICHODERINAE [Shattuck (1992c) gave a generic revision]

- *103. *Anonychomyrma* Donisthorpe 1947 AU, OR Shattuck, 1992c
104. *Bothriomyrmex* Emery 1869 AU, OR, PA
105. *Dolichoderus* Lund 1831 AU, NA, NT, OR, PA Shattuck, 1992c
Hypoclinea Mayr 1855 =Dolichoderus
Ireneae Donisthorpe 1938 =Polyrhachis
(FORMICINAE)
106. *Iridomyrmex* Mayr 1862 AU, OR Shattuck, 1992a
Leptomyrme Mayr 1862 →[AU]
Liometopum Mayr 1861 →[NA, NT, PA]
Monoceratoclinea Wheeler 1935 =Dolichoderus
Myrmapatetes Wheeler 1929 =Odontomachus
(PONERINAE)
- *107. **Loweriella** Shattuck 1992c OR
- *108. *Ochetellus* Shattuck 1992 AU, OR Shattuck, 1992a
- *109. *Philidris* Shattuck 1992 AU, OR Shattuck, 1992a
Semonius Forel 1910 =Tapinoma
110. *Tapinoma* Foerster 1855 AU, ET, MA, NA, NT,
OR, PA

111. *Technomyrmex* Mayr 1872 AU, ET, MA, OR, PA
Trumeria Forel 1895 → [AU]
Zatapinoma Wheeler 1928 =*Tapinoma*

FORMICINAE

- Acantholepis* Mayr 1861 =*Baroniurbania*
112. *Acropyga* Roger 1862 AU, ET, NA, NT, OR, PA
Andragathus Emery 1922 =*Paratrechina* [Agosti & Bolton, 1990]
Anacantholepis Santschi 1914 =*Plagiolipsis*
113. *Anoplolepis* Santschi 1914 AU, ET, OR
- *114. *Baroniurbania* Pagliano & Scaramozzino 1989 AU, ET, OR, PA
- *115. *Bregmatomyrma* Wheeler 1929 OR
- *116. *Calomyrmex* Emery 1895 AU, OR
117. *Camponotus* Mayr 1861 AU, ET, MA, NA, NT, OR, PA [Subg. *Colobopsis* is sometimes treated as genus]
118. *Cataglyphis* Foerster 1850 ET, OR, PA Agosti, 1990
Chapmanella Wheeler 1930 =*Euprenolepis*
119. *Cladomyrma* Wheeler 1920 OR Agosti, 1991
120. *Echinopla* F.Smith 1857 AU, OR
- *121. *Euprenolepis* Emery 1906 AU, OR
- *122. ***Forelophilus*** Kutter 1931 OR
123. *Formica* Linnaeus 1758 NA, NT, OR, PA
124. ***Gesomyrmex*** Mayr 1868 OR
Hemioptica Roger 1962 =*Polyrhachis*
125. *Lasius* Fabricius 1804 NA, OR, PA Wilson, 1955
Mesozena F. Smith 1860 =*Echinopla*
Myrmecocystus Wasmal 1838 → [NA]
126. ***Myrmoterus*** Forel 1893 OR Moffett, 1985
Nylanderia Emery 1906 =*Paratrechina*
127. *Oecophylla* F.Smith 1860 AU, ET, OR
128. *Opisthopis* Emery 1893 AU, OR
- *129. ***Overbeckia*** Viehmeyer 1915 OR
130. *Paratrechina* Motschulsky 1863 AU, ET, MA, NA, NT, OR, PA Trager, 1984
131. *Plagiolipsis* Mayr 1861 AU, ET, MA, OR, PA

132. Prenolepis Mayr 1861 NA, NT, OR, PA
Polyergus Latreille 1804 →[NA, PA]
133. Polyrhachis F.Smith 1857 AU, ET, OR, PA Hung, 1970
134. **Pseudaphomomyrmex** Wheeler OR
 1920
135. Pseudolasius Emery 1886 AU, ET, OR

References

- Agosti, D., 1990. Review and classification of *Cataglyphis* (Hymenoptera, Formicidae). J. Nat. Hist. 24:1457-1505.
- Agosti, D., 1991. Revision of the oriental ant genus *Cladomyrma*, with an outline of the higher classification of the Formicinae (Hymenoptera, Formicidae). Syst. Ent. 16: 293-310.
- Agosti, D. & B. Bolton, 1990. The identity of *Andragathus*, a forgotten formicine ant genus (Hym., Formicidae). Ent. Mon. Mag. 126:75-77.
- Baroni Urbani, C., 1975a. Contributo alla conoscenza dei generi *Belonopelta* Mayr e *Leiopelta* gen. n. (Hymenoptera: Formicidae). Mit. Schweiz. Entomol. Ges. 48:295-310.
- Baroni Urbani, C., 1975b. Primi reperi del genere *Calyptomyrmex* Emery nel subcontinente Indiano. Ent. Basil. 1: 395-411.
- Baroni Urbani, C., 1977a. Ergebniss der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel. Hymenoptera: Fam. Formicidae Genus *Myriella*. Entom. Basiliensia 2: 411-414.
- Baroni Urbani, C., 1977b. Ergebniss der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel. Hymenoptera: Fam. Formicidae Genus *Stenamma* con una nuova specie del Kashmir. Entom. Basiliensia 2: 415-422.
- Baroni Urbani, C., 1977c. Materiali per una revisione della sottofamiglia Leptanillinae Emery (Hymenoptera: Formicidae). Entom. Basiliensia 2: 427-488.
- Bolton, B., 1974. A revision of the Palearctic arboreal ant genus *Cataulacus* F. Smith (Hymenoptera: Formicidae). Bull. Brit. Mus. Nat. Hist. (Ent.) 30:1-105.
- Bolton, B., 1976. The ant tribe Tetramoriini (Hymenoptera: Formicidae): constituent genera, review of smaller genera and revision of *Triglyphothrix* Forel. Bull. Brit. Mus. Nat. Hist. (Ent.) 34:281-378.
- Bolton, B., 1977. The ant tribe Tetramoriini (Hymenoptera: Formicidae): the genus *Tetramorium* Mayr in the Oriental and Indo-Australian regions, and in Australia. Bull. Brit. Mus. Nat. Hist. (Ent.) 38:67-151.
- Bolton, B., 1981. A revision of the ant genera *Meranoptus* F.Smith, *Dicroaspis* Emery and *Calyptomyrmex* Emery (Hymenoptera: Formicidae) in the Ethiopian zoogeographical region. Bull. Brit. Mus. Nat. Hist. (Ent.) 42: 43-81.
- Bolton, B., 1982. Afrotropical species of the myrmicine ant genera *Cardiocondyla*, *Leptothorax*, *Melissotarsus*, *Messor* and *Cataulacus* (Formicidae). Bull. Brit. Mus. Nat. Hist. (Ent.) 45: 307-370.
- Bolton, B., 1983. The Afrotropical dacetine ants (Formicidae). Bull. Brit. Mus. Nat. Hist. (Ent.) 46: 267-416.
- Bolton, B., 1986. A taxonomic review of the tetramorine ant genus *Rhoptromyrmex* (Hymenoptera: Formicidae). Syst. Ent. 11:1-17.
- Bolton, B., 1987. A review of the *Solenopsis* genus-group and revision of Afrotropical *Monomorium* Mayr (Hymenoptera: Formicidae). Bull. Brit. Mus. Nat. Hist. (Ent.) 54:263-452.
- Bolton, B., 1988a. A review of *Paratpula* Wheeler, a forgotten genus of myrmicine ant (Hym., Formicidae). Ent. Mon. Mag. 124:125-143.
- Bolton, B., 1988b. *Secostruma*, a new subterranean tetramorine ant genus (Hymenoptera: Formicidae). Syst.

- Ent. 13:263-270.
- Bolton, B., 1990a. Abdominal characters and status of the cerapachyine ants (Hymenoptera: Formicidae). Jour. Nat. Hist. 24: 53-68.
- Bolton, B., 1990b. The higher classification of the ant subfamily Leptanillinae (Hymenoptera: Formicidae). Syst. Ent. 15: 267-282.
- Bolton, B., 1990c. Army ants reassessed: the phylogeny and classification of the doryline section (Hymenoptera: Formicidae). Jour. Nat. Hist. 24: 1339-1364.
- Brown, W. L., Jr., 1955. Ant taxonomy. E. L. Kessel, ed. A century of progress in the natural sciences, 1853-1953, pp. 569-572. California Academy of Sciences, San Francisco.
- Brown, W. L., Jr., 1958. Contributions toward a reclassification of the Formicidae. II. Tribe Ectatommini (Hymenoptera). Bull. Mus. Comp. Zool. Harv. 118:175-362.
- Brown, W. L., Jr., 1960. Contributions toward a reclassification of the Formicidae. III. Tribe Amblyoponini (Hymenoptera). Bull. Mus. Comp. Zool. Harv. 122:144-230.
- Brown, W. L., Jr., 1964. Genus *Rhaptomyrmex*, revision and key to species. Pilot Register of Zoology, cards 11-19.
- Brown, W. L., Jr., 1973. A comparison of the Hylean and Congo-West African rain forest ant faunas. B.J. Meggers, E.S. Ayensu and W.D. Duckworth eds., Tropical forest ecosystems in Africa and South America: a comparative review, pp. 161-185. Smithsonian Institution Press, Washington, D.C.
- Brown, W. L., Jr., 1975. Contributions toward a reclassification of the Formicidae. V. Ponerinae, tribes Platythreini, Cerapachyini, Cylandromyrmecini, Acanthostichini and Aenictogitini. Search - Agr. Entom. 5:1-115.
- Brown, W. L., Jr., 1976. Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section A. Introduction, subtribal characters, genus *Odontomachus*. Stud. Ent. 19:17-171.
- Brown, W. L., Jr., 1978. Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section B. Genus *Anochetus* and bibliography. Stud. Ent. 20:549-638.
- Chapman, J. W. & S. R. Capco, 1951. Check list of the ants (Hymenoptera: Formicidae) of Asia. Monographs of the Institute of Science and Technology, Philippines, no.1, 327 pp.
- Emery, C., 1921-22. Hymenoptera, family Formicidae, subfamily Myrmicinae. P. Wytsman ed. Genera Insectorum, fasc. 174. Louis Desmet-Verteneuil, Brussels. 397 pp.
- Ettershank, G., 1966. A generic revision of world Myrmicinae related to *Solenopsis* and *Pheidologeton* (Hymenoptera: Formicidae). Aust. J. Zool. 14: 73-171.
- Holldobler, B. & E. O. Wilson, 1990. The ants. The Belknap Press of Harvard University Press, Cambridge. 732 pp.
- Hung, A. C. F., 1970. A revision of ants of the subgenus *Polyrhachis* Fr. Smith (Hymenoptera: Formicidae: Formicinae). Orient. Ins. 4: 1-36.
- Imai, H. T., W.L. Brown, Jr., M. Kubota, H.-S. Yong & Y. P. Tho, 1983. Chromosome observation on tropical ants from Western Malaysia. II. Ann. Rep. Nat. Inst. Genetics (Japan) (34):66-69.
- Imai, H. T. & M. Kubota, 1981. Karyological survey of Indian ants. Some preliminary reports. Nucleus (India), 24: 93-96.
- Imai, H. T., M. Kubota, W.L. Brown, Jr., M. Ihara, M. Tohari & R.I. Pranata, 1985. Chromosome observation on tropical ants from Indonesia. Ann. Rep. Nat. Inst. Genetics (Japan) (35):46-48.
- Lincoln R.J. & G.A. Boxshall, 1987. The Cambridge illustrated dictionary of natural history, Cambridge University Press. 413 pp.
- Moffett, M. W., 1985. Revision of the genus *Myrmoteras* (Hymenoptera: Formicidae). Bull. Mus. Comp. Zool. Harv. 151:1-53.
- Myrmecologists Society (Japan) (ed.), 1988. A list of the ants of Japan with common Japanese names. Tokyo, 50

- pp. (In Japanese).
- Myrmecological Society of Japan (ed.), 1989. A guide for the identification of Japanese ants (I). Ponerinae, Cerapachyinae, Pseudomyrmecinae, Dorylinae and Leptanillinae (Hymenoptera: Formicidae). Tokyo, 42 pp. (In Japanese).
- Myrmecological Society of Japan (ed.), 1991. A guide for the identification of Japanese ants (II). Dolichoderinae and Formicinae (Hymenoptera: Formicidae). Tokyo, 56 pp. (In Japanese).
- Myrmecological Society of Japan (ed.), 1992. A guide for the identification of Japanese ants (III). Myrmicinae and supplement to Leptanillinae (Hymenoptera: Formicidae). Tokyo, 94 pp. (In Japanese).
- Ogata, K., 1991. Ants of the genus *Myrmecia* Fabricius: a review of the species groups and their phylogenetic relationships (Hymenoptera: Formicidae: Myrmecinae). Syst. Ent. 16: 353-381.
- Shattuck, S. O., 1992a. Review of the dolichoderinae ant genus *Iridomyrmex* Mayr with descriptions of three new genera (Hymenoptera: Formicidae). J. Aust. Entomol. Soc. 31: 13-18.
- Shattuck, S. O., 1992b. Higher classification of the ant subfamilies Aneuretinae, Dolichoderinae and Formicinae (Hymenoptera: Formicidae). Syst. Entomol. 17: 199-206.
- Shattuck, S. O., 1992c. Generic revision of the ant subfamily Dolichoderinae (Hymenoptera: Formicidae). Sociobiol. 21: 1-181.
- Taylor, R. W., 1985. The ants of the Papuanian genus *Dacetonops* (Hymenoptera: Formicidae: Myrmicinae). G. E. Ball ed., Taxonomy, phylogeny and zoogeography of beetles and ants, pp.41-67. Dordrecht.
- Taylor, R. W., 1983. Descriptive taxonomy: past, present, and future. E. Highley & R. W. Taylor eds., Australian systematic entomology: a bicentenary perspective, pp.93-134. CSIRO, Melbourne.
- Taylor, R. W., 1990a. New Asian ants of the tribe Basicerotini, with an on-line computer interactive key to the twenty-six known Indo-Australian species (Hymenoptera: Formicidae: Myrmicinae). Invertebr. Taxon. 4: 397-425.
- Taylor, R. W., 1990b. Notes on the ant genera *Romblonella* and *Willowskiella*, with comments on their affinities, and the first descriptions of Australian species (Hymenoptera: Formicidae: Myrmicinae). Psyche, 97: 281-296.
- Taylor, R. W., 1991. Nomenclature and distribution of some Australian ants of the Myrmicinae (Hymenoptera: Formicidae). Mem. Queensland Mus. 30: 599-614.
- Tjan, K. N., H. T. Imai, M. Kubota, W. L. Brown Jr., W. H. Gotwald Jr., H. S. Yong & C. Leh., 1986. Chromosomal observations of Sarawak ants. Ann. Rep. Nat. Inst. Genetics (Japan) (36): 57-58.
- Ward, P. S., 1980. A systematic revision of the *Rhytidoponera impressa* group in Australia and New Guinea. Aust. J. Zool. 28: 475-498.
- Ward, P. S., 1990. The ant subfamily Pseudomyrmecinae (Hymenoptera: Formicidae): generic revision and relationship to other formicids. Syst. Ent. 15: 1449-459.
- Wilson, E. O., 1959. Adaptive shift and dispersal in a tropical ant fauna. Evolution, 13: 122-144.
- Wilson, E. O., 1964. The true army ants of the Indo-Australian area (Hymenoptera: Formicidae: Dorylinae). Pac. Ins. 6: 427-483.
- Wilson, E. O., 1987. The current status of ant taxonomy. J. C. Trager ed., Advances in Myrmecology, pp. 3-10. New York.
- Wilson, E. O., 1988. The current status of biological diversity. E.O. Wilson ed., Biodiversity, pp. 3-18. National Academy Press, Washington, D.C.
- Wilson, E. O. & W. L. Brown, Jr., 1956. New parasitic ants of the genus *Kygidris*, with notes on the ecology and behavior. Ins. Soc. 3: 439-454.